

Lifetime prevalence of non-melanoma and melanoma skin cancer in Australian recreational and competitive surfers

Climstein, Mike; Furness, James; Hing, Wayne; Walsh, Joe

Published in:
Photodermatology Photoimmunology and Photomedicine

DOI:
[10.1111/phpp.12247](https://doi.org/10.1111/phpp.12247)

Licence:
Unspecified

[Link to output in Bond University research repository.](#)

Recommended citation(APA):
Climstein, M., Furness, J., Hing, W., & Walsh, J. (2016). Lifetime prevalence of non-melanoma and melanoma skin cancer in Australian recreational and competitive surfers. *Photodermatology Photoimmunology and Photomedicine*, 32(4), 207-213. <https://doi.org/10.1111/phpp.12247>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.

1 **Lifetime prevalence of non-melanoma and melanoma skin cancer in Australian**
2 **recreational and competitive surfers**

3

Abstract

Background/purpose

Surfing is one of the most popular outdoor aquatic activities in Australia with an estimated 2.7 million recreational surfers however, Australia has long been recognized as having the highest incidence of melanoma in the world, and it is the most common type of cancer in young Australians. The aim of this study was to investigate the lifetime prevalence of non-melanoma (basal cell carcinoma (BCC), squamous cell carcinoma (SCC)), and melanoma skin cancers in Australian recreational and competitive surfers.

Methods

Australian surfers were invited to complete an online surveillance survey to determine the lifetime prevalence of non-melanoma and melanoma skin cancers.

Results

A total 1,348 surfers (56.9% recreational) participated in this study, of which 184 surfers reported a skin cancer (competitive n=96, recreational n=87). Of non-melanoma and melanoma cancers reported, BCC was the most common (6.8%), followed by melanoma (1.4%) and SCC (0.6%). The relative risk was higher ($p<0.001$) in competitive versus recreational surfers (OR 1.74 (CI 1.28-2.31)). There was a higher ($p<0.05$) number of skin cancers reported on the face (23.5%), back (16.4%) and arms (12.4%). There were significant trends ($p<0.001$) in reported skin cancers between competitive and recreational surfers, as well as significantly ($p<0.001$) more skin cancers reported in males (14.6%) than females (9.4%).

Conclusion

26 Based upon these findings, individuals who surf are advised to regularly utilize sun protection
27 strategies (avoid peak ultra violet radiation (10am-3pm), rashvest, hat and sunscreen) and
28 primary care physicians are recommended to regularly screen their patients who surf.

29

30 **Keywords:** physical activity; public health; questionnaire survey; sun exposure; skin
31 neoplasms; ultraviolet radiation

32

33

Introduction

Surfing is one of the most popular outdoor aquatic activities in Australia with an estimated 2.7 million recreational surfers (1). As a result the expected risk of skin cancer in surfers due to long periods of exposure to ultraviolet radiation (UVR) (2), wearing less clothing and reflection from water is of great concern. According to the Cancer Council, nearly one third of all Australians had a skin cancer, making it the most common type of cancer in Australians (3). Non-melanoma skin cancers (NMSC) were the most prevalent with over 750,000 treated each year. Australia has also been recognized as having the highest incidence of melanoma skin cancer (MSC) in the world (4-6), and it is the most common cancer in Australians aged 15 to 29 years (7), although this has recently been challenged (8). Melanoma is also the third most common form of skin cancer, the mechanism of which is attributed to sunlight exposure, particularly UVR within the 250 to 400 nm wavelength (9, 10). It is estimated in 2015 there will be an estimated 938,000 treatments related to skin cancer in Australia, reportedly costing the medical benefits scheme \$703 million for diagnosis, treatment and pathology (11). A recent report on the epidemiological incidence of keratinocyte skin cancer revealed basal cell carcinoma (BCC) was the most common type of skin cancer in the Australian population followed by squamous cell carcinoma (SCC) (12), the report emphasized UVR exposure is of critical importance in Australian as a mechanism for the development of skin cancer. Despite the popularity of surfing there is very limited research (13) available on the prevalence of skin cancers in surfers. Dozier and Wagner (12) reported the results of a free skin cancer screening conducted by dermatologists in Texas (USA) surfers (n=49). The majority (73.4%) of the surfers screened were identified as having atypical moles (36.7%), actinic keratosis (40.8%) and BCCs (16.3%). Significance ($p < 0.47$) was only identified with regard to BCCs in the surfers as opposed to a self-selected general population (16.3% versus 3.2%, respectively). There were no lesions identified that were suggestive of SCCs or melanomas. The purpose

of this study was to investigate the lifetime prevalence of non-melanoma and melanoma skin cancers in recreational and competitive surfers in Australia.

Methods

Study design

This research utilized a commercially available, customizable cloud-based survey in a cross-sectional observational study design.

Sample

Given the wide geographic distribution of surfing in Australia, an on-line survey was developed to allow for national data collection on surfing injuries (14) and determine the lifetime prevalence of non-melanoma and melanoma skin cancers in recreational and competitive surfers. We defined lifetime prevalence as the proportion of the population who, at some point in their lifetime up to the time of this survey, has ever had a melanoma (MSC) or non-melanoma skin cancer (NMSC). Media (broadcast, print and radio) promotion and direct contact with national surfing organizations and surfing clubs (n=103) was utilized to promote the study to all individuals who were currently surfing either recreationally or competitively in Australia. Ethics approval for this study was granted by Bond University Human Research Ethics committee (RO1540).

Survey

The survey consisted of two primary sections. Section 1 contained questions that pertained to participants physiological demographics (such as age, height, mass) and their surfing specific

demographics (board type) and surfing exposure (average hours per week and weeks per year for the previous year). Additionally, we inquired if the participants surfed during peak UVR which is recognized as 10am to 3pm, inclusive (15, 16). We also queried the participants surf status (recreational or competitive) and if the latter whether it was for their local club, international or the World Championship Tour. Section 2 pertained to surfing injuries (14) (acute and chronic) and included questions pertaining to the lifetime prevalence of non-melanoma and melanoma skin cancers. Participants were instructed to report only NMSCs and MSCs by type and location that have only been diagnosed and/or treated by either a general practitioner (GP) or dermatologist. This question specified that diagnosis needed to be confirmed by tissue biopsy, however participants were not required to upload either medical reports of diagnosed skin cancers or associated pathology reports. As most of the participants would be unaware of their Fitzpatrick skin phototype (17) we simplified skin types to the following; fair, medium, olive, brown and black (18).

Survey questions consisted of array, single choice, multiple choice, list dropdown, numerical input and short answer free text. Filters (where appropriate) were utilized to ensure participants provided electronic informed consent (individuals under the age of 18 years were required to have parental or guardian supervision whilst completing the survey), met entry criteria, expedite completion of the survey and help ensure responses were within logical limits. Additionally, participants were informed they could only complete one survey per participant and the online survey mechanism allowed each participant to only complete the survey once. All participants were required to provide informed consent electronically to enable access to the survey which was active for six months.

Statistical analysis

Data normality was assessed by investigating kurtosis, skewness, Q-Q plots, as well as the Kolmogorov-Smirnov (KS) test with Lilliefors significance correction. Heteroscedasticity was assessed using Levene's test. Statistical analyses were completed using the IBM Statistical Package for the Social Sciences (SPSS, version 22) and included demographics (mean \pm SD), correlations, t-tests and chi-square tests to determine significance of differences between groups, alpha was set at $p < 0.05$ *a priori*.

Results

Sample characteristics

Table 1 depicts the physiological and surfing demographics of all participants. A total of 1,348 surfers completed the survey, the majority of which were male (91.3%) and surfing recreationally (56.9%). The overall mean age of participants was 35.8 years (range 11-70) with competitive surfers ($n=581$, 43.1%) significantly ($p < 0.05$) older (+3.9%) and slightly shorter (-0.05%) than recreational surfers. With regard to surfing demographics competitive surfers reported surfing significantly ($p < 0.001$) more hours per week (+37.2%), weeks per year (+13.4%) and total hours per year (+43.8%). There was no difference ($p > 0.05$) with regard to surfing exposure between genders. The majority of surfers rode short boards (82%), followed by longboards (10.9%) and mini-mals (7.1%)

Insert Table 1 approximately here

Lifetime prevalence of skin cancer

A total of 184 of the survey participants (13.6%) reported being diagnosed and/or treated for a NMSC and/or MSC (Figure 1), there were a greater number of skin cancers reported in competitive (Table 2) than recreational surfers ($p<0.05$). Competitive surfers had reported proportionally more skin cancers than recreational surfers (16.5% and 11.4%, respectively). The relative risk of developing a skin cancer was significantly ($p<0.001$) higher in competitive versus recreational surfers (odds ratio 1.74 (CI 1.28-2.31)). An additional 37 participants (2.7%) reported having a skin cancer however, were unable to recall the specific type. There was a trend ($p=0.06$) for competitive surfers to surf more often during peak UVR as opposed to recreational surfers (50.9% versus 39.0%, respectively).

Insert Figure 1 approximately here

Regarding location, NSW participants reported the highest incidence of skin cancers (38.3%) followed closely by Queensland (37.8%). Other states included Victoria (8.5%), followed closely by Western Australia (8.0%), then South Australia (4.8%) and Tasmania (0.5%).

With regard to the total cohort, BCC's were the most frequently (6.8%) reported NMSC followed by MSCs (1.4%) and NMSC SCC's (0.6%). Of note is the number of participants who reported multiple types of skin cancers; BCC and SCC (1.7%), BCC and SCC and melanoma (0.3%) and BCC and melanoma (0.2%). Competitive surfers reported more BCC's (+19.6%) however, surprisingly there were more MSC reported by recreational surfers (Table 2). There were no significant differences between the type of skin cancer and surfing status (recreational versus competitive). The NMSC BCC was primarily reported in NSW ($n=51$) followed closely by Queensland ($n=47$), this was similar to SCCs (NSW, $n=17$; Queensland,

n=12). Melanoma skin cancers were most prevalent in Queensland (n=11), followed closely by NSW (n=9).

Insert Table 2 approximately here

Skin cancer by site

With respect to the site of the skin cancers (Figure 2), there was a significantly higher incidence of skin cancers on the face ($p<0.001$), back ($p<0.001$) and arms ($p<0.05$) whereas the feet, thigh and neck had the lowest number of reported skin cancers. Although our cohort of female surf participants is the largest dataset of lifetime prevalence of skin cancer to date, for comparative purposes this study was limited due to the relatively small number (n=117) of female participants in comparison to male participants (n=1231). The location of MSC was most commonly reported on the back (27.8%) followed by the face (16.3%) and shoulder and arms (11.4%).

Insert Figure 2 approximately here

Key demographical and physiological risk factors

There was a negative correlation between age and time spent surfing with older surfers surfing less hours per week ($p<0.0001$) and less weeks per year ($p<0.005$). As would be expected, however important to confirm, older surfers had also surfed for more years than their younger counterparts ($r=0.73$, $p<0.0001$). Those surfers who surfed more hours per week, tended to also surf more weeks of the year ($p<0.0001$).

Surfers reporting a skin cancer had higher body mass index (BMI) ($p < 0.0001$), older in age ($p < 0.0001$), and mass ($p < 0.0001$), though years surfing was identified as the most significant factor ($p < 0.0001$).

There was a significant difference between genders as males reported a higher number of skin cancers versus females (0.36 vs. 0.12, $p < 0.0001$). This was however, unsurprising as males also had surfed for more years than females ($p < 0.0001$) and this was shown to also be related to the number of skin cancers reported.

Insert Figure 1 here

Competitive surfers were found to have a significantly ($p < 0.001$) higher proportion of skin cancers than recreational surfers however, competitive surfers had surfed longer ($p < 0.0001$) and were older (36.6 vs. 35.2yrs).

Skin cancer and skin color

With regard to skin colour, there was a significant ($p < 0.005$) relationship between the number of skin cancers and lighter skin as fair and medium skin participants reported the highest frequency (43.4% and 46.7%, respectively) of NMSCs (86.4%) and MSCs (13.6%). However, it should be noted that the limited number of brown ($n=1$) and black ($n=0$) skinned participants is not a representative sample.

Discussion

This study represents the largest survey that has investigated NMSCs and MSC in both recreational and competitive surfers. Our findings are in agreement with the only previously published study (13) which reported a similar (16.3%) cohort prevalence of skin cancer. Although that study did not report SCC per se, the authors did report a high prevalence (40.8%) of actinic keratosis which has been shown to give rise to SCCs in 44 to 82% of cases (19). The prevalence of BCC in our study is less than previously reported (13) (16.3%) despite our study investigating lifetime prevalence versus incidence.

The risk of skin cancer development has also been investigated in other outdoor recreational and sporting groups and shown to be an independent risk factor for BCC (2). Lichte and colleagues (20) investigated the incidence of skin cancer in mountain guides and reported a similar incidence of BCCs (7.1%) and a slightly higher incidence of SCCs (1.4%). The cohort (n=283) of adult males only identified 1 MSC. Skin cancer has also been investigated in outdoor farm workers (21), the proportionate MSC incidence (0.18) was found to be higher than urinary bladder cancer (0.59) however, lower than kidney cancer (1.60), liver cancer (4.24), prostate cancer (1.13) and uterine cancer (2.08).

Our data analysis demonstrated that the total number of years surfing was a more significant factor with regard to skin cancer than current surfing activity (hours per week). However, this may be mitigated by the fact that surfers who were older surfed less than their younger counterparts. A consideration is that if older surfers have always surfed less than younger counterparts (an assumption that may not be valid), the extent of skin cancers reported in future may be far higher once the currently younger cohort of surfers have spent an equivalent number of years exposed to UVR whilst surfing.

220

221 With regard to the tendency ($p>0.05$) for less skin cancers on the top of head, face, ears and
222 neck in females, this was possibly attributed to the protective factor of longer hair however,
223 this was not evaluated in the survey and is only speculative.

224

225 These findings are in agreement with previous research which have reported a lower incidence
226 of skin cancers in races and ethnicities with darker skin (22, 23), as these races and ethnicities
227 have been shown to have a higher melanin content (22). Bradford (24) previously reported in
228 the United States Caucasians had the highest incidence of skin cancers (35-40%) followed by
229 Hispanics (4-5%) Asians (2-4%) and Afro-Americans (1-2%).

230

231 This study also found a lifetime prevalence for melanoma to be 1.4% within a surfing
232 population whereas a 27 year prevalence of melanoma in Australia was found to be 0.6% in
233 2007 (25), comparatively the rate in the USA (26) was reported to be considerably higher (3.8%)
234 and is expected to rise significantly in the near future (27). This comparison needs to be viewed
235 cautiously as timeframes differ; however as this is the only comparable Australian dataset it
236 highlights the potential threat of melanoma in patients who surf.

237

238 Non-melanoma skin cancers are the most common cancer in Australia and the highest in the
239 world (11). These cancers also place a high financial burden on the Australian health care
240 system with an estimated yearly cost of \$700 million. Given the large proportion of
241 Australians who surf and numbers of reported NMSC and MSC within this cohort, surfers may
242 account for a notable proportion of all skin cancers within Australia. The authors acknowledge
243 that the generalizability of our results is limited to Australian surfers.

244

245 With regard to limitations to this study, selection bias may have existed as individuals without
 246 internet access would not have had the opportunity to have participated, we attempted to
 247 minimize information bias by requiring melanoma and non-melanoma data to only be entered
 248 when diagnosed and/or treated by the participants GP or specialist. We believe there was no
 249 confounding bias as we did not investigate casual relationships. A further limitation to this
 250 study was the exclusion of investigating sun protection strategies which have been shown to
 251 be effective in reducing the risk of skin cancer in other outdoor athletes (28).

252

253 **Conclusions**

254 Surfers have previously been surveyed successfully (29-31) and we believe this study adds
 255 insight into the lifetime prevalence of non-melanoma and melanoma skin cancers in the
 256 Australian surfing cohort. Although the data obtained was self-report which has recently come
 257 under question (32), participants were required to only report skin cancers diagnosed and/or
 258 treated by their GP or medical specialist. Given this study's findings with respect to the lifetime
 259 prevalence of skin cancer in Australian surfers, GP's are recommended to rigorously screen
 260 their patients who surf as based upon our findings they appear to be at increased risk of
 261 developing either a non-melanoma and/or melanoma skin cancer. Screenings are important
 262 preventative measures as GP screening examinations has previously been shown to be effective
 263 in reducing premature death from skin cancers (5). Furthermore, GP's should also advise
 264 patients to utilize multiple sun protection strategies specific to surfing (avoid peak UVR,
 265 rashvest, hat and sunscreen (9, 33, 34)). Previous investigators (35) have shown that sun safety
 266 education programs are effective in changing sun protective behaviours, particularly for non-
 267 melanoma and melanoma skin cancers.

268 Additionally, lighter skinned surfers with an extensive surfing history appear to a key subgroup
269 at greatest risk for skin cancer. The key high incidence areas of exposure need recognition and
270 strategies to reduce the incidence of skin cancers in these areas would be worthy of
271 consideration (i.e. the face, back and arms).

272

273

274 **Acknowledgements**

275 The authors wish to thank Surfing Australia's High Performance Centre, Surfing Australia,
276 Surfing Queensland, SurfRider Foundation, Tracks magazine, Swellnet.com.au, Surfing Life,
277 Fitnessnetwork.com, NBN TV, Nine News, ABC radio (Gold Coast, Sunshine Coast),
278 Sunday Mail, and the Gold Coast Bulletin for assisting in the promotion of this research.

279

280 All authors contributed to the design of the research study, Dr. Furness performed the
281 research, all authors contributed to the statistical analyses and writing of this paper.

282

283 **Disclosure and competing interests**

284 None declared

285

REFERENCES

1. Stark A. Surfing Australia Annual Report. 2013.
2. Moehrle M. Outdoor sports and skin cancer. *Clin Dermatol*. 2008; 26:12-15.
3. Cancer Council of Australia. Skin Cancer Statistics and Issues. 2015.
4. Stanton W. Primary prevention of skin cancer in Australia: Report of the Sun Protection Programs Working Party. 1996.
5. Sladden MJ, Ward JE, Del Mar CB, Lowe JB. Skin cancer screening by Australian family physicians: variation with physician beliefs and geographic locality. *Am J Prev Med*. 1999; 17:142-146.
6. Volkov A, Dobbinson S, Wakefield M, Slevin T. Seven-year trends in sun protection and sunburn among Australian adolescents and adults. *Aust N Z J Public Health*. 2013; 37:63-69.
7. Perez D, Kite J, Dunlop SM, Cust AE, Goumas C, Cotter T, et al. Exposure to the 'Dark Side of Tanning' skin cancer prevention mass media campaign and its association with tanning attitudes in New South Wales, Australia. *Health Educ Res*. 2015.
8. Olsen CM, Williams PF, Whiteman DC. Turning the tide? Changes in treatment rates for keratinocyte cancers in Australia 2000 through 2011. *J Am Acad Dermatol*. 2014; 71:21-26 e21.
9. Moehrle M. Outdoor sports and skin cancer. *Clin Dermatol*. 2008; 26:12-15.
10. Kanavy HE GM. Ultraviolet radiation and melanoma. *Semin Cutan Med Surg*. 2011; 30:222-228.
11. Fransen M, Karahalios A, Sharma N, English DR, Giles GG, Sinclair RD. Non-melanoma skin cancer in Australia. *Med J Aust*. 2012; 197:565-568.
12. Refshauge A. Cancer Survival and Prevalence in Australia: Period estimates from 1982 to 2010. 2012.
13. Dozier S, Wagner RF, Jr., Black SA, Terracina J. Beachfront screening for skin cancer in Texas gulf coast surfers. *South Med J*. 1997; 90:55-58.
14. Furness JW, Hing WA, Abbott A, Walsh J, Climstein M. Retrospective analysis of chronic injuries in recreational and competitive surfers: Injury location, type and mechanism. *International Journal of Aquatic Research and Education*. 2014; 8:228-287.
15. Agency ARPANS. Radiation Protection. 2014.
16. Gonzaga ER. Role of UV light in photodamage, skin aging, and skin cancer: importance of photoprotection. *Am J Clin Dermatol*. 2009; 10 (Suppl 1):19-24.
17. Eilers S, Bach DQ, Gaber R, Blatt H, Guevara Y, Nitsche K, et al. Accuracy of self-report in assessing Fitzpatrick skin phototypes I through VI. *JAMA dermatology*. 2013; 149:1289-1294.
18. Jablonski N. Human Evolutionary Biology. In: Muehlenbein MP, ed. *Human Evolutionary Biology*. England: Cambridge University Press, 2010: 192-213.
19. Cohen JL. Actinic keratosis treatment as a key component of preventive strategies for nonmelanoma skin cancer. *The Journal of Clinical and Aesthetic Dermatology*. 2010; 3:39-44.
20. Lichte V, Dennenmoser B, Dietz K, Hafner HM, Schlagenhauff B, Garbe C, et al. Professional risk for skin cancer development in male mountain guides--a cross-sectional study. *J Eur Acad Dermatol Venereol*. 2010; 24:797-804.
21. Mills PK, Shah P. Cancer incidence in California farm workers, 1988-2010. *Am J Ind Med*. 2014; 57:737-747.
22. Brenner M, Hearing VJ. The protective role of melanin against UV damage in human skin. *Photochem Photobiol*. 2008; 84:539-549.
23. Control CfD. Skin Cancer Rates by Race and Ethnicity. 2015.
24. Bradford PT. Skin cancer in skin of color. *Dermatol Nurs*. 2009; 21:170-177, 206; quiz 178.
25. Refshauge A, Kalisch D. Cancer survival prevalence in Australia. 2012.
26. Siegal R, Ma J, Zou Z, Jemal A. Cancer statistics, 2014. 2015.

27. Weir HK, Thompson TD, Soman A, Moller B, Leadbetter S. The past, present, and future of cancer incidence in the United States: 1975 through 2020. *Cancer*. 2015.
28. Noble-Jerks J, Weatherby RP, Meir R. Self-reported skin cancer protection strategies and location of skin cancer in retired cricketers: a case study from membership of the Emu Cricket Club. *J Sci Med Sport*. 2006; 9:441-445.
29. Howden J, Danks J, McCluskey P, Gillett M, Ghabrial R. Surfboard-related eye injuries in New South Wales: a 1-year prospective study. *Med J Aust*. 2014; 201:532-534.
30. Furness J, Hing W, Abbott A, Walsh J, Sheppard JM, Climstein M. Retrospective Analysis of Chronic Injuries in Recreational and Competitive Surfers: Injury Location, Type, and Mechanism. *International Journal of Aquatic Research and Education*. 2014; 8:277-287.
31. Furness J, Hing W, Walsh J, Abbott A, Sheppard JM, Climstein M. Acute Injuries in Recreational and Competitive Surfers Incidence, Severity, Location, Type, and Mechanism. *The American journal of sports medicine*. 2015; 43:1246-1254.
32. Holm AS, Wulf HC. Self-reported skin cancer is unreliable. *Eur J Epidemiol*. 2015; 30:159-162.
33. Glanz K, Mayer JA. Reducing ultraviolet radiation exposure to prevent skin cancer methodology and measurement. *Am J Prev Med*. 2005; 29:131-142.
34. Saraiya M. Recommendations to prevent skin cancer by reducing exposure to ultraviolet radiation. *Am J Prev Med*. 2004; 27:467-470.
35. Jinna S, Adams BB. Ultraviolet radiation and the athlete: risk, sun safety, and barriers to implementation of protective strategies. *Sports Med*. 2013; 43:531-537.

Table 1. Descriptive statistics for participants (values are mean (\pm SD)). 95% Confidence intervals (95% CI).

Variable	Cohort (n=1,348)	Recreational (n=767)	Competitive (n=581)	P value
Age (yrs)	35.84 (13.08)	35.2 (12.2)	36.7 (14.1)	.04
Weight (kg)	78.59 (12.79)	78.3 (11.9)	78.9 (13.7)	NS
Height (cm)	178.16 (9.03)	178.6 (9.1)	177.7 (8.6)	NS
BMI (kg/m ²)	24.73 (3.75)	24.5 (3.3)	24.9 (3.5)	NS
Surfing demographics				
Surfing:				
• Hours/week	6.7 (5.6)	5.4 (4.4)	8.6 (6.6)	.001
• Weeks/year	40.5 (13.6)	38.0 (14.6)	43.8 (11.4)	.001
• Hours/year	305.5 (291.2)	228.7 (214.3)	406.9 (343.7)	.001

Table 2. Lifetime incidence of non-melanoma and melanoma skin cancers in participants
(number, percentage)

Type	Cohort (n=1,348)	Recreational (n=767)	Competitive (n=581)	P value
BCC (n)	123	56 (45.5%)	67 (54.5%)	NS
Melanoma (n)	25	15 (60.0%)	10 (40.0%)	NS
SCC (n)	36	17 (47.2%)	19 (52.8%)	NS
Total (n)	184	88 (47.8%)	96 (52.2%)	NS